



Geographic variation in pigmentation patterns of Indo-Pacific humpback dolphins (*Sousa chinensis*) in Chinese waters

BINGYAO CHEN, THOMAS A. JEFFERSON, LIN WANG, HUILI GAO, HONGKE ZHANG, YU ZHOU, XINRONG XU, AND GUANG YANG*

Jiangsu Key Laboratory for Biodiversity and Biotechnology, Nanjing Normal University, 1 Wenyuan Road, Nanjing 210023, China (BC, LW, HG, XX, GY)

Clymene Enterprises, Lakeside, CA 92040, USA (TAJ)

National Hepu Dugong Nature Reserve administration station, 49 Gongyuan Road, Beihai 536000, China (HZ, YZ)

* Correspondent: 08255@njnu.edu.cn

The pigmentation patterns of Indo-Pacific humpback dolphins have not been well studied throughout most of the species' range. In the present study, both the subjective scoring method and a newly developed quantified scoring method were employed to evaluate the spotting intensity of 137 humpback dolphin individuals from the Xiamen and Beibu Gulf putative populations, including the Dafengjiang-Nanliujiang River Estuary (DRE) and Shatian-Caotan (SC) communities. Both scoring methods indicated that spotting intensity on the dorsal fin was lower than that on the body of humpback dolphins in all 3 groups. The SC and DRE humpback dolphins had significantly greater dorsal fin, body spotting intensity, and greater differences between spotting on the dorsal fin and body than Xiamen dolphins, while no differences were found between SC and DRE. The pigmentation variation is related to age class, young dolphins have more spotting density than adults, and young and adults showed similar geographical variation as above. The present paper, in combination with previous research, clarifies the general pattern of pigmentation for Chinese humpback dolphins. Eastern Taiwan Strait and Pearl River Estuary populations represent 2 extreme patterns of pigmentation, while the Xiamen population, SC community, and DRE community seem to be intermediate. The results suggest that these groups should be viewed as demographically distinct forms.

在大多数分布区内，中华白海豚的色素沉着模式尚未得到很好的研究。本研究应用两种方法：前人应用的主观评分法和新开发的数字化评分法，来评估厦门、南流江-大风江河口（DRE），和沙田-草潭（SC）群体中137头中华白海豚个体的斑点密度。两种评分方法均表明，3个群的中华白海豚背鳍的斑点密度低于身体的斑点强度。SC和DRE中华白海豚的背鳍斑点强度、身体斑点强度，和背鳍和身体上的斑点差值，均显著高于厦门中华白海豚，而SC和DRE之间没有显著差异。色素沉着与年龄相关，我们的研究表明年轻和成年个体呈现出与上述相似的结果。本文结合前人的研究，阐明了中国水域内中华白海豚色素沉着的普遍规律：东台湾海峡和珠江河口的种群分别代表了两种极端的色素沉着模式，而厦门种群，北部湾种群（包括SC社群和DRE社群）代表中间形式，这些群体是统计学显著不同的存在类型。

Key words: Beibu Gulf, coloration, Indo-Pacific humpback dolphin, photo-identification, pigmentation pattern, Xiamen

Compared to most mammals, cetaceans are not a particularly colorful group (Stockin and Visser 2005), but geographical variation of color patterns does occur in cetaceans (Perrin 1972). Despite limited light penetration in the oceans, coloration patterns are important in cetaceans, presumably having ecological function and adaptive value (Perrin 2009; Caro et al. 2011). Most species show rather limited variation within their typical pigmentation patterns (Stockin and Visser 2005); however, a

few species or genera show extensive color variation, for example, beluga whales (*Delphinapterus leucas*—Brodie 1989) and humpback dolphins (*Sousa* spp.—Jefferson and Karczmarski 2001; Wang et al. 2008; Jefferson and Rosenbaum 2014).

In dolphins of the genus *Sousa*, young calves and juveniles are typically devoid of spots in all 4 species (Jefferson and Rosenbaum 2014; Brown et al. 2015; Jefferson et al. 2015). Adults of the Indo-Pacific humpback dolphin (*Sousa*

chinensis, hereafter humpback dolphin) have much more extensive white coloration on the body than the other 3 species of *Sousa* (Jefferson and Rosenbaum 2014; Jefferson et al. 2015). The base pigmentation of adult humpback dolphins is white, although they often have small dark spots on the body or dorsal fin (Jefferson and Karczmarski 2001; Wang et al. 2008).

Within 8 putative Chinese humpback dolphin populations (distributed from east to west: Ningde—Chen et al. 2012; Eastern Taiwan Strait [ETS]—Wang et al. 2004, 2007; Xiamen—Liu and Huang 2000; Chen et al. 2008, 2011; Chou et al. 2013; Shantou—Wu 2010; Pearl River Estuary [PRE] including Hong Kong—Jefferson 2000; Hung 2008; Zhanjiang—Xu et al. 2012, 2015; Sanya—Dong et al. 2017; and the northern Beibu Gulf including Dafengjiang-Nanliujiang River Estuary [DRE] and Shatian-Caotan [SC] groups—Chen et al. 2009, 2016; Fig. 1), 3 populations were sampled to investigate pigmentation patterns (Wang et al. 2008). The results indicated that dorsal fins of ETS humpback dolphins were more spotted when compared to those of Xiamen and PRE populations (Wang et al. 2008), and this formed part of the evidence defining the ETS population as a new subspecies of Indo-Pacific humpback dolphin (*S. c. taiwanensis*—Wang et al. 2015). For other populations, however, the pigmentation patterns have not been extensively researched and remained largely undocumented.

In this paper, we selected the Xiamen population, which could be compared with a previous study, and the westernmost, most distant groups from Xiamen (SC and DRE) for determining geographic variation. We used the same method as Wang et al. (2008) to subjectively score pigmentation of identified humpback dolphins. In addition, a quantitative method was developed to evaluate the spotting on the dorsal fin and

the body of humpback dolphins. This paper provides updated information on the geographic variation of pigmentation patterns within Chinese waters.

MATERIALS AND METHODS

The Xiamen, DRE, and SC Indo-Pacific humpback dolphins were photographed in the wild. The research team had a special permit from Animal Research Ethics Committee of Nanjing Normal University. During encounters with dolphins, efforts were made to photograph all members of the group of dolphins seen during a sighting and to avoid disturbance. If dolphins displayed boat avoidance behavior, such as deep dives, long down times, changes in direction, or fleeing and fast swimming, encounters were ended.

The DRE humpback dolphins and SC dolphins possibly belong to the same population inhabiting the northern Beibu Gulf during our research. Because the photographic comparison found no matches of individuals between DRE and SC, they were considered as separate communities in this paper. Their data were therefore analyzed separately. In total, photographs of 137 individual dolphins were selected for comparison: Xiamen ($n = 36$), SC ($n = 23$), and DRE ($n = 78$).

Spotting was scored by 2 methods: 1) a subjective scoring method developed by Wang et al. (2008, 2015), and 2) a quantified scoring method developed by us. For the subjective scoring method, the density of spotting on the dorsal fin and on the body below the dorsal fin was scored on a scale from 1 to 4 (Wang et al. 2008; see examples in Fig. 2, left). The photographs were scored independently by 6 scorers. Four scorers were cetacean researchers (each familiar with this species), and 2 were researchers with no experience with humpback dolphins.




Fig. 1.—The distribution of putative Indo-Pacific humpback dolphin (*Sousa chinensis*) “populations” in China. ETS = Eastern Taiwan Strait population; PRE = Pearl River Estuary population; SC = Shatian-Caotan community; DRE = Dafengjiang-Nanliujiang River Estuary community.

The subjective scoring method resulted in integer score data, which possibly produced experience-related and subjective biases. Therefore, we developed a new quantified scoring method, which is more objective. The threshold method was applied to digitize the photographs of dolphins into white background and black spots (Fig. 2, right). Then, the percentages of black spots on the dorsal fin and on the body were calculated using Image-Pro Plus 6.0 (Ahern 1998). The differences of body-to-dorsal fin spotting (body score minus dorsal fin score) were also calculated.

The differences of spotting scores of humpback dolphins from different groups were tested. One-way analyses of variance (ANOVAs) and multiple pairwise *t*-tests were used to test between populations (Xiamen, DRE, and SC), based on the scores of dorsal fin, body, and the difference of body-to-dorsal fin. A 1-sample *t*-test was conducted comparing the pigmentation scores of Xiamen, DRE, or SC with the mean scores of ETS or PRE from data in Wang et al. (2008; Table 1). All the tests were performed with a 0.05 significance level, using IBM SPSS Statistics version 19 (Gray and Kinnear 2012). For the subjective score scatterplot, the data (subjective body and

dorsal fin score) for ETS and PRE were preliminarily estimated from figure 2 of Wang et al. (2015).

To explore the influence of age class and consider sample size, we re-classified all dolphins into 2 main age classes (young: mottled and speckled, adult: spotted and unspotted). Then, we compared differences in their pigmentation pattern.

RESULTS

Based on the subjective scoring method, the mean intensity of spotting on dorsal fins of all 3 groups was less than that on the bodies (Table 1), i.e., Xiamen (2.03 versus 2.74), SC (2.97 versus 3.34), and DRE (2.76 versus 3.16). The mean difference of body-to-dorsal fin of Indo-Pacific humpback dolphins of the 3 populations varied from 0.37 to 1.40, but was mostly < 1.0 (Fig. 3). Xiamen had significantly less spotting intensity on the dorsal fin and body than SC, and had significantly less spot intensity on the dorsal fin and greater score difference of body-to-dorsal fin (Tables 1 and 2). The frequency distribution of different spotting intensity of the dorsal fin of Xiamen humpback dolphins was significantly different from that of SC or DRE groups, whereas the frequency distributions for the body and




Fig. 2.—The reference for subjective scoring (left) and quantified scoring methods (right) of spotting on the dorsal fins and bodies of Indo-Pacific humpback dolphins (*Sousa chinensis*). In subjective scoring, Score 1 = little to no spotting; Score 2 = light spotting; Score 3 = moderate spotting; Score 4 = heavy spotting. Photos by B. Chen and X. Xu.

Table 1.—Mean subjective scores of spotting intensity of Indo-Pacific humpback dolphins (*Sousa chinensis*) in Chinese waters. ETS = Eastern Taiwan Strait; PRE = Pearl River Estuary; SC = Shatian-Caotan; DRE = Dafengjiang-Nanliujiang River Estuary.

Population	<i>n</i>	Body score (mean ± SE)	Dorsal fin score (mean ± SE)	Difference of body-to-dorsal fin scores (mean ± SE)	Reference
ETS	31	2.92 ± 0.13	3.3 ± 0.13	−0.38 ± 0.08	Wang et al. (2008)
Xiamen	10	2.6 ± 0.37	1.53 ± 0.14	1.07 ± 0.24	Wang et al. (2008)
Xiamen	36 ^a	2.74 ± 0.15	2.03 ± 0.14	0.72 ± 0.09	This study
PRE	186	2.66 ± 0.07	1.43 ± 0.04	1.23 ± 0.05	Wang et al. (2008)
SC	23	3.34 ± 0.14	2.97 ± 0.19	0.37 ± 0.11	This study
DRE	78	3.16 ± 0.10	2.76 ± 0.11	0.40 ± 0.06	This study

^a1 adult (a black patch in the left front body) was the same individual used in Wang et al. (2008).




Fig. 3.—The frequency distributions of subjective scores (left) and quantified scores (right) of spotting intensity of the dorsal fin (top), body (middle), and body-to-dorsal fin comparison (bottom) of Indo-Pacific humpback dolphins (*Sousa chinensis*). SC = Shatian-Caotan population; DRE = Dafengjiang-Nanliujiang River Estuary population.

body-to-dorsal fin difference (Fig. 3, left) among the 3 groups were similar.

Based on the quantified scoring method, the mean intensity of spotting on dorsal fins of all 3 groups was less than that on the bodies (Table 1), which is the same pattern detected by the subjective scoring results. The mean spotting percentage on the dorsal fins and on the bodies of humpback dolphins was greatest

in DRE and was lowest in Xiamen (Table 3). The difference of body-to-dorsal fin was the lowest in Xiamen and the highest in SC. The frequency distributions of spotting intensity of the dorsal fin and difference of body-to-dorsal fin were similar among the 3 groups, whereas it was different in spotting of the body (Fig. 3, right). When comparing the pigmentation percentage of the 3 dolphin groups, the SC and DRE humpback dolphins had

Table 2.—Matrix of significant test results of the subjective scoring of spotting intensity of Indo-Pacific humpback dolphins (*Sousa chinensis*) from 5 putative populations in Chinese waters. Data above (right) and below (left) of the diagonal are associated *P*-values and the difference between both tested populations, respectively. ETS = Eastern Taiwan Strait; PRE = Pearl River Estuary; SC = Shatian-Caotan; DRE = Dafengjiang-Nanliujiang River Estuary.

		ETS	PRE	Xiamen	SC	DRE
ETS	Body				< 0.05	
	Dorsal fin		< 0.001 ^a	< 0.001		< 0.05
PRE	Difference		< 0.001 ^a	< 0.001	< 0.001	< 0.001
	Body				< 0.001	< 0.05
	Dorsal fin	1.77 ^a		< 0.05	< 0.001	< 0.001
Xiamen	Difference	1.45 ^a		< 0.001	< 0.001	< 0.001
	Body				< 0.05	
	Dorsal fin	-1.387	0.598		< 0.05	< 0.05
SC	Difference	1.145	-0.513			< 0.05
	Body	0.419	0.679	0.595		
	Dorsal fin		1.535	0.937		
DRE	Difference	0.754	-0.856			
	Body		0.504			
	Dorsal fin	-0.396	1.332	0.734		
	Difference	0.658	-0.827	-0.314		

^aData from Wang et al. (2008).

Table 3.—The quantified scores of spotting intensity of Indo-Pacific humpback dolphins (*Sousa chinensis*) in Chinese waters. SC = Shatian-Caotan; DRE = Dafengjiang-Nanliujiang River Estuary. Superscript letters in the same row (a, b) identify significant pairwise differences.

		Xiamen (n = 36)	SC (n = 23)	DRE (n = 78)
Body score (%)	Mean	26.58 ^{a,b}	46.83 ^a	43.86 ^b
	Low	0.28	0.37	0.08
	High	75.55	86.87	91.13
Dorsal fin score (%)	Mean	11.22 ^{a,b}	29.66 ^a	24.56 ^b
	Low	0.74	0	0
	High	47.87	74.74	84.54
Difference of body-to-dorsal fin scores (%)	Mean	15.35	17.17	20.96
	Low	-5.1	-6.98	-5.58
	High	55.14	56.36	84.39

Table 4.—The subjective and quantified scores of spotting intensity of young (mottled and speckled dolphins) and adults (unspotted and spotted dolphins) Indo-Pacific humpback dolphins (*Sousa chinensis*) in Chinese waters.

	Population	Age	n	Body score	Dorsal fin score	Difference of body-to-dorsal fin scores
Quantified score	Xiamen	Young	15	47.79%	18.30%	29.49%
	SC		9	69.25%	57.83%	13.14%
	DRE		32	71.72%	45.77%	26.34%
	Xiamen	Adult	21	11.42%	6.17%	6.06%
	SC		14	32.43%	11.56%	20.87%
	DRE		46	22.72%	8.60%	14.32%
Subjective score	Xiamen	Young	15	3.51	2.60	0.91
	SC		9	3.87	3.82	0.04
	DRE		32	3.89	3.48	0.41
	Xiamen	Adult	21	2.20	1.62	0.58
	SC		14	3.00	2.41	0.59
	DRE		46	2.66	2.26	0.40

significantly greater dorsal fin and body spotting intensity than Xiamen dolphins, while no distinctive difference in pigmentation was found between SC and DRE groups (Table 4).

The scatterplot of the quantified scores indicated that SC and DRE humpback dolphins were separated from those in Xiamen (Fig. 4, top left). In the trend line analysis, the following

functions were calculated: Xiamen: $Y = 0.3651X + 0.0152$; SC: $Y = 0.8006X - 0.0783$; and DRE: $Y = 0.6726X - 0.0494$. The slope for Xiamen was significantly shallower than that for SC and DRE, and the intercept for Xiamen was positive, whereas the intercepts for SC and DRE were negative. These points indicated that the spotting intensity of Xiamen humpback




Fig. 4.—Scatterplot of general and age-related subjective scores (left) and quantified scores (right) of spotting intensity of dorsal fins, bodies, and difference between body and dorsal fin of 3 Indo-Pacific humpback dolphin (*Sousa chinensis*) populations and 2 communities. The middle and bottom figures show the age class variation. The data for ETS and PRE in the left figure were obtained from figure 2 of Wang et al. (2015). ETS = Eastern Taiwan Strait; PRE = Pearl River Estuary; SC = Shatian-Caotan; DRE = Dafengjiang-Nanliujiang River Estuary.

dolphins was different from that of SC and DRE communities. The young dolphins in SC showed differences from Xiamen and DRE (middle left), but the adults (bottom left) were similar to the general trend (top left).

When comparing all 6 geographic areas from our data plus those of Wang et al. (2008, 2015), including ETS and PRE

groups, the scatterplot of subjective score and trend lines showed differences in pigmentation among 5 groups (Fig. 4, top right). The trend line equations produced were: ETS: $Y = 0.8315X + 0.9393$; Xiamen: $Y = 0.7659X + 0.073$; PRE: $Y = 0.4577X + 0.2335$; SC: $Y = 1.0525X - 0.5493$; and DRE: $Y = 0.8481X + 0.078$. The slopes varied from 0.4577 in PRE to

1.0525 in SC. The intercept of ETS (0.9393) was significantly larger than that in all other populations, and the intercept of SC (−0.5493) was the smallest. The young (middle right) and adult (bottom right) dolphins showed patterns similar with the general trend (top right).

For mottled–speckled (young) and unspotted–spotted adults, significant geographical pigmentation differences were found (Table 4). For the quantified score of adults, the density of body spotting ($F_{2,80} = 4.928$, $P = 0.01$; multiple comparisons: Xiamen–SC, $P = 0.026$; Xiamen–DRE, $P = 0.011$) and differences of body-to-dorsal fin ($F_{2,80} = 4.786$, $P = 0.011$; Xiamen–SC, $P = 0.014$; Xiamen–DRE, $P = 0.019$) were geographically significant. The young dolphins showed geographical differences in pigmentation on dorsal fins ($F_{2,54} = 13.516$, $P < 0.001$; Xiamen–DRE, $P < 0.001$; Xiamen–SC, $P < 0.001$), body ($F_{2,54} = 12.12$, $P < 0.001$; Xiamen–DRE, $P = 0.001$; Xiamen–SC, $P = 0.007$), and difference of body-to-dorsal fin ($F_{2,54} = 3.875$, $P = 0.027$; DRE–SC, $P = 0.017$; SC–Xiamen, $P = 0.013$).

For subjective scores, the adult dolphins showed geographical variation in pigmentation on dorsal fins ($F_{2,80} = 6.224$, $P = 0.003$; Xiamen–DRE, $P = 0.006$; Xiamen–SC, $P = 0.008$) and body ($F_{2,80} = 4.521$, $P = 0.014$; Xiamen–SC, $P = 0.006$). Young dolphins also showed differences in pigmentation on dorsal fins geographically ($F_{2,54} = 17.346$, $P < 0.001$; Xiamen–DRE, $P = 0.001$; Xiamen–SC, $P < 0.001$; SC–DRE, $P = 0.038$), body ($F_{2,54} = 8.051$, $P = 0.001$; Xiamen–DRE, $P = 0.029$), and difference of body-to-dorsal fin ($F_{2,54} = 8.405$, $P < 0.001$; Xiamen–SC, $P = 0.001$; SC–DRE, $P = 0.001$).

DISCUSSION

In this study, we used only high-quality photos for analyses. Only photos with sharp focus were analyzed, and any blurred photos or photos with poor lighting were excluded. Although pigmentation changes over time, the change is limited throughout a 6- to 8-year period. Therefore, for each dolphin, only a single clear photograph was selected, and we believe this is adequate.

Both subjective and quantified scoring methods produced similar results, and we believe they are both appropriate methods. For the Xiamen population, the scores in the present paper were different from those of Wang et al. (2008), although the general pattern was similar. This might have resulted from different target dolphins, sample sizes, or age structures. Individual variation in scoring methods may have also played a part. The photos of the Xiamen animals used in the Wang et al. (2008, 2015) study were taken in the late 1990s ($n = 10$), whereas ours were taken in 2007–2010 ($n = 36$). Fortunately, we photographed the same individual dolphin (Btop in figure 2 of Wang et al. 2008). The pigmentation of this dolphin changed very little over at least 9 years.

Ontogenetic changes in coloration are widespread in cetaceans, such as those found in killer whales (*Orcinus orca*—Evans et al. 1982), humpback whales (*Megaptera novaeangliae*—Rosenbaum et al. 1995), pantropical spotted dolphins

(*Stenella attenuata*—Perrin and Hohn 1994), and Indo-Pacific bottlenose dolphins (*Tursiops aduncus*—Bichell et al. 2018), and geographic differences aid in taxonomic studies and the identification of various geographic forms, subspecies, and species (see Perrin 2009).

Of the 6 studied populations, the ETS and PRE groups seemed to represent 2 extreme pigmentation patterns, while Xiamen, SC, and DRE were likely to show more intermediate patterns (Tables 1 and 2; Fig. 4). Significant differences were found among these populations, and this suggests that there is distinctive geographical variation. Further, the differentiation of pigmentation patterns in the ETS population was used as one of the critical features for describing that form as a new subspecies, *S. c. taiwanensis* (Wang et al. 2015). The variation in pigmentation of these populations is irregular along their geographic locations (east to west, or south to north). Analyses of DNA supported the irregular geographic variation (Chen et al. 2010). The unique sample from the Beibu Gulf shared the same haplotype (CH02) with dolphins in Xiamen (Chen et al. 2010). Further research is needed to understand how DNA evolution is related to the change of pigmentation.

Pigmentation and spotting intensities in *S. chinensis* are known to be related to age and sex (Jefferson et al. 2012; Jefferson and Rosenbaum 2014), like the situation in some other small cetaceans, e.g., *S. attenuata* and *Tursiops* sp. (Krzyszczuk and Mann 2012). Despite the age-related variation in pigmentation, young dolphins (mottled and speckled) and adults (spotted and unspotted) both showed similar patterns of geographic variation of pigmentation. However, effects of sex remain unclear, due to the challenge of sex identification in the wild.

Combined, the present paper and previous work by Wang et al. (2008, 2015) have analyzed 4 of 8 putative Indo-Pacific humpback dolphin “populations” in China. There is general agreement among these studies, and they each reflect the general framework of variation in pigmentation of Indo-Pacific humpback dolphins in Chinese waters. For the other 4 populations, similar research to that presented here should be done. However, Ningde and Shantou appear to have small populations, and few identified individuals are currently available (e.g., there are only 4 dolphins from Ningde in our database). This would make such research challenging. However, due to its importance, we suggest that this be pursued in the future, despite the difficulties.

ACKNOWLEDGMENTS

We thank the National Key Programme of Research and Development, Ministry of Science and Technology (2016YFC0503200), the National Natural Science Foundation of China (31630071), NSF of Jiangsu Province of China (BK20131400), the Priority Academic Program Development of Jiangsu Higher Education Institutions (PAPD), Ocean Park Conservation Foundation Hong Kong, OPCFHK (MM01-1213, MM01-1314), Major project of hydrobios resources in Jiangsu province, and Hepu Dugong National Nature Reserve

Management Station for funding. We also thank P. A. Olson for her subjective scoring and suggestions on an early version of the manuscript, and the anonymous reviewers.

LITERATURE CITED

- AHERN, K. 1998. Image-Pro Plus 6.0 (Media Cybernetics Company, MD, USA) Image-Pro Plus. Biotechnology Software and Internet Journal 15:10–15.
- BICHELL, L. M. V., E. KRZYSZCZYK, E. M. PATTERSON, AND J. MANN. 2018. The reliability of pigment pattern-based identification of wild bottlenose dolphins. *Marine Mammal Science* 34:113–134.
- BRODIE, P. F. 1989. The white whale *Delphinapterus leucas* (Pallas, 1776). Pp. 119–144 in *Handbook of marine mammals* (S. H. Ridgway and R. R. Harrison, eds.). Academic Press, London, United Kingdom.
- BROWN, A. M., ET AL. 2015. Sexual dimorphism and geographic variation in dorsal fin features of Australian humpback dolphins, *Sousa sahulensis*. *Advances in Marine Biology* 73:273–314.
- CARO, T., T. K. BEEMAN, T. STANKOWICH, AND H. WHITEHEAD. 2011. The functional significance of colouration in cetaceans. *Evolutionary Ecology* 25:1231–1245.
- CHEN L., S. CABALLERO, K. ZHOU, AND G. YANG. 2010. Molecular phylogenetics and population structure of *Sousa chinensis* in Chinese waters inferred from mitochondrial control region sequences. *Biochemical Systematics and Ecology* 38:897–905.
- CHEN, B., D. ZHENG, J. JU, X. XU, K. ZHOU, AND G. YANG. 2011. Range patterns of resident Indo-Pacific Humpback dolphins (*Sousa chinensis*, Osbeck 1765) in Xiamen, China: implications for conservation and management. *Zoological Studies* 50:751–762.
- CHEN, B., D. ZHENG, L. WANG, X. XU, AND G. YANG. 2012. The northernmost distribution of Indo-Pacific Humpback dolphin (*Sousa chinensis*) in the world: evidence from preliminary survey in Ningde, China. *Pakistan Journal of Zoology* 44:1209–1214.
- CHEN, B., D. ZHENG, G. YANG, X. XU, AND K. ZHOU. 2009. Distribution and conservation of the Indo-Pacific humpback dolphin in China. *Integrative Zoology* 4:240–247.
- CHEN, B., ET AL. 2008. Abundance, distribution and conservation of Chinese white dolphins (*Sousa chinensis*) in Xiamen, China. *Mammalian Biology* 73:156–164.
- CHEN, B., ET AL. 2016. Conservation status of the Indo-Pacific humpback dolphin (*Sousa chinensis*) in the northern Beibu Gulf, China. *Advances in Marine Biology* 73:119–139.
- CHOU, L. S., W. L. CHANG, Y. J. WU, AND H. Y. YU. 2013. Indo-Pacific humpback dolphins in the waters of Kinman: distribution, habitat usage and population structure. *Journal of National Park* 23:1–10 (in Chinese).
- DONG, L., M. LIU, J. DONG, AND S. LI. 2017. Acoustic occurrence detection of a newly recorded Indo-Pacific humpback dolphin population in waters southwest of Hainan Island, China. *The Journal of the Acoustical Society of America* 142:3198–3204.
- EVANS, W. E., A. V. YABLOKOV, AND A. E. BOWLES. 1982. Geographic variation in the color pattern of killer whales (*Orcinus orca*). *Report of the International Whaling Commission* 32:687–694.
- GRAY, C. D., AND P. R. KINNEAR. 2012. IBM SPSS statistics 19 made simple. Psychology Press, New York.
- HUNG, S. K. 2008. Habitat use of Indo-Pacific humpback dolphins (*Sousa chinensis*) in Hong Kong. Ph.D. dissertation, University of Hong Kong, Hong Kong, China.
- JEFFERSON, T. A. 2000. Population biology of the Indo-Pacific humpbacked dolphin in Hong Kong waters. *Wildlife Monographs* 144:1–65.
- JEFFERSON, T. A., S. K. HUNG, K. M. ROBERTSON, AND F. I. ARCHER. 2012. Life history of the Indo-Pacific humpback dolphin in the Pearl River Estuary, southern China. *Marine Mammal Science* 28:84–104.
- JEFFERSON, T. A., AND L. KARCZMARSKI. 2001. *Sousa chinensis*. *Mammalian Species* 655:1–9.
- JEFFERSON, T. A., AND H. C. ROSENBAUM. 2014. Taxonomic revision of the humpback dolphins (*Sousa* spp.), and description of a new species from Australia. *Marine Mammal Science* 30:1494–1541.
- JEFFERSON, T. A., M. A. WEBBER, AND R. L. PITMAN. 2015. *Marine mammals of the world: a comprehensive guide to their identification*. 2nd ed. Academic Press, San Diego, California.
- KRZYSZCZYK, E., AND J. MANN. 2012. Why become speckled? Ontogeny and function of speckling in Shark Bay bottlenose dolphins (*Tursiops* sp.). *Marine Mammal Science* 28:295–307.
- LIU, W. H., AND Z. G. HUANG. 2000. Distribution and abundance of Chinese white dolphin (*Sousa chinensis*) in Xiamen. *Acta Oceanologica Sinica* 22:95–101 (in Chinese).
- PERRIN, W. F. 1972. Color patterns of spinner porpoises (*Stenella* cf. *S. longirostris*) of the eastern Pacific and Hawaii, with comments on delphinid pigmentation. *Fishery Bulletin* 70:983–1003.
- PERRIN, W. F. 2009. Coloration. Pp. 243–255 in *Encyclopedia of marine mammals* (W. F. Perrin, B. Würsig, and H. Thewissen, eds.). Academic Press, Amsterdam, The Netherlands.
- PERRIN, W. F., AND A. A. HOHN. 1994. Pantropical spotted dolphin *Stenella attenuata*. Pp. 173–191 in *Handbook of marine mammals*. Volume 5. The first book of dolphins (S. H. Ridgway and R. J. Harrison, eds.). Academic Press, Inc., London, United Kingdom.
- ROSENBAUM, H. C., ET AL. 1995. Geographic variation in ventral fluke pigmentation of humpback whale *Megaptera novaeangliae* populations worldwide. *Marine Ecology Progress Series* 124:1–7.
- STOCKIN, K. A., AND I. N. VISSER. 2005. Anomalous pigmented common dolphins (*Delphinus* sp.) off northern New Zealand. *Aquatic Mammals* 31:43–51.
- WANG, J. Y., S. K. HUNG, AND S. C. YANG. 2004. Records of Indo-Pacific humpback dolphins, *Sousa chinensis* (Osbeck, 1765): from the waters of western Taiwan. *Aquatic Mammals* 30:189–196.
- WANG, J. Y., S. K. HUNG, S. C. YANG, T. A. JEFFERSON, AND E. R. SECCHI. 2008. Population differences in the pigmentation of Indo-Pacific humpback dolphins, *Sousa chinensis*, in Chinese waters. *Mammalia* 72:302–308.
- WANG, J. Y., S. C. YANG, AND S. K. HUNG. 2015. Diagnosability and description of a new subspecies of Indo-Pacific humpback dolphin, *Sousa chinensis* (Osbeck, 1765), from the Taiwan Strait. *Zoological Studies* 54:36.
- WANG, J. Y., S. C. YANG, S. K. HUNG, AND T. A. JEFFERSON. 2007. Distribution, abundance and conservation status of the Eastern Taiwan Strait population of Indo-Pacific humpback dolphins, *Sousa chinensis*. *Mammalia* 71:157–165.
- WU, Y. 2010. The survival status of Chinese white dolphin in Guangdong Province, China. Ph.D. dissertation, Shantou University, Shantou, Guangdong Province, China.
- XU, X., J. SONG, Z. ZHANG, G. YANG, AND K. ZHOU. 2015. The world's second largest population of humpback dolphins in the waters of Zhanjiang deserves the highest conservation priority. *Scientific Reports* 5:8147.
- XU, X., Z. ZHANG, L. MA, G. YANG, AND K. ZHOU. 2012. Site fidelity and association patterns of Indo-Pacific humpback dolphins off the east coast of Zhanjiang, China. *Acta Theriologica* 57:99–109.

Submitted 23 October 2017. Accepted 12 June 2018.

Associate Editor was Rochelle Constantine.